

Table 22.9  
Cancer of the Pancreas (Invasive)  
SEER<sup>a</sup> Relative Survival (Percent)  
By Year of Diagnosis

All Races, Males and Females

Survival Time	Year of Diagnosis																			
	1975- 1979	1980- 1984	1985- 1989	1990- 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
1-year	14.2	15.9	17.1	18.5	17.7	19.1	20.0	19.6	22.2	21.9	23.8	25.2	22.6	23.1	26.9	25.9	27.9	29.9	29.4	30.1
2-year	5.8	6.3	6.8	8.5	8.2	9.3	9.9	8.7	10.5	10.9	11.4	11.9	11.2	11.9	12.8	13.0	13.7	14.9	15.3	
3-year	3.6	4.1	4.7	5.8	5.2	6.3	7.1	5.6	6.5	7.3	7.3	8.4	7.4	8.3	8.8	9.2	9.6	10.2		
4-year	2.9	3.3	3.8	4.8	3.9	4.8	5.5	4.5	5.6	6.0	6.1	6.8	5.6	6.4	7.0	7.9	8.1			
5-year	2.5	2.8	3.3	4.2	3.6	4.2	5.0	3.8	5.1	5.1	5.0	6.2	4.8	5.5	6.1	7.3				
6-year	2.3	2.5	2.9	4.0	3.2	3.8	4.5	3.7	4.6	4.6	4.7	5.7	4.3	5.0	5.6					
7-year	2.1	2.3	2.7	3.6	3.1	3.5	4.3	3.4	4.3	4.2	4.1	5.1	4.0	4.7						
8-year	1.9	2.1	2.5	3.4	2.8	3.4	4.2	3.1	3.9	3.7	3.9	4.8	3.8							
9-year	1.8	2.0	2.2	3.2	2.7	3.3	4.0	2.8	3.8	3.6	3.8	4.7								
10-year	1.7	1.9	2.2	2.9	2.6	3.2	3.9	2.7	3.6	3.5	3.4									
11-year	1.7	1.8	2.1	2.8	2.5	2.9	3.8	2.5	3.3	3.5										
12-year	1.7	1.7	2.1	2.7	2.3	2.8	3.6	2.4	3.1											
13-year	1.6	1.6	2.0	2.6	2.2	2.6	3.3	2.3												
14-year	1.5	1.6	1.9	2.5	1.8	2.6	3.1													
15-year	1.5	1.5	1.8	2.3	1.7	2.6														
16-year	1.5	1.4	1.7	2.2	1.7															
17-year	1.4	1.4	1.7	2.2																
18-year	1.3	1.3	1.7																	
19-year	1.2	1.3	1.6																	
20-year	1.2	1.3	1.6																	

<sup>a</sup> Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).